

The involvement of Fc gamma receptor gene polymorphisms in Kawasaki disease

M. Biezeveld,^{*,†} J. Geissler,[†]

M. Merkus,[‡] I. M. Kuipers,^{*}

J. Ottenkamp^{*,§} and T. Kuijpers^{*,†}

**Emma Children's Hospital, Academic Medical Centre (AMC), University of Amsterdam, Amsterdam, the Netherlands, †Sanquin Research at CLB, University of Amsterdam, the Netherlands, ‡Center for Pediatric Clinical Epidemiology (KEK), Emma Children's Hospital, Amsterdam, the Netherlands, and §Leiden University Medical Center (LUMC), Pediatric Cardiology, Leiden, the Netherlands*

Accepted for publication 26 October 2006

Correspondence: Maarten Biezeveld, Emma Children's Hospital, Academic Medical Centre (AMC), Room G8-205, University of Amsterdam, Meibergdreef 9, 1105 AZ, Amsterdam, the Netherlands.

E-mail: m.h.biezeveld@amc.uva.nl

Summary

Kawasaki disease is an acute febrile syndrome in infancy, characterized by vasculitis of medium-sized arteries. Without treatment the disease can lead to coronary artery lesions (CAL) in approximately 25% of the children. Therapy consists of intravenous immunoglobulins (IVIG), leading to a decrease of complications to 5–16%. Little is known about the working mechanisms of IVIG. In this study we evaluated the involvement of Fc γ receptors (Fc γ Rs) in Kawasaki disease by the determination of the frequency of known single nucleotide polymorphisms (SNPs) in the genes coding for the Fc γ Rs and compared this with frequencies in a cohort of healthy controls. There was no difference in the distribution of the functionally relevant genotypes for Fc γ RIIa-131H/R, Fc γ RIIb-232I/T, Fc γ RIIIa-158 V/F and Fc γ RIIIb-NA1/NA2 between the patient group and the healthy controls. Furthermore, there were no polymorphisms linked to the disease severity as indicated by the absence or development of CAL during the disease. Altered transcription or expression of Fc γ R on specific cell types of the immune system may still play a role in susceptibility and treatment success, but at a level different from the functional SNPs in Fc γ R genes tested in this study.

Keywords: Fc gamma receptors, intravenous immunoglobulin, Kawasaki disease, single nucleotide polymorphism

Introduction

Kawasaki disease is an acute febrile syndrome in infancy, which is characterized by vasculitis of mainly the medium-sized arteries. Because there is no specific test to diagnose the disease, diagnosis is made on clinical criteria [1,2].

The incidence of Kawasaki disease is 5–17 per 100 000 in Europe and the United States, mainly in children under the age of 5 years. For unknown reasons the incidence in Japan is 135 per 100 000 [3]. In approximately 25% of the children vasculitis will lead to coronary artery lesions (CAL) as detected by echocardiography, showing this to be the leading cause of acquired heart disease in children.

Therapy consists of a single high dose of intravenous immunoglobulins (IVIG, 2 g/kg) infused in 8–12 h and aspirin. This therapy decreases complications to 5–16% [4,5]. To date, little is still known about the working mechanisms of IVIG. One of the proposed mechanisms of action of IVIG depends on the functional interaction with Fc γ receptors (Fc γ Rs). Fc γ Rs are receptors for the constant region of immunoglobulins (the Fc region) [6]. They play an important role in the immune response against invading

microorganisms. Fc γ Rs enhance the identification and destruction of noxious material, foreign as well as endogenous, by opsonization. Hereby they form the link between the humoral and cellular parts of the immune system, as well as a crucial link between innate and adaptive immunity [7].

The group of Fc γ Rs comprises three different classes: Fc γ RI (CD64), Fc γ RII (CD32) and Fc γ RIII (CD16). Depending on their expression on effector cells, Fc γ R exert different effects. For example, on phagocytes they mediate phagocytosis, endocytosis, antibody-dependent cellular cytotoxicity (ADCC) and induction of respiratory burst [6]. Apart from expression, a 100-fold difference in IgG-binding affinity exists. Fc γ RI is considered as high-affinity receptor and types II and III [8,9] as low-affinity receptors. Of these receptors, Fc γ RII is the most widely distributed class and is expressed on most types of blood cells [6,10]. Fc γ RII is the only Fc γ R member with its own signalling motif and contains, depending on the isoform, either an immunoreceptor tyrosine-based activation motif (ITAM) or an immunoreceptor tyrosine-based inhibitory motif (ITIM). Fc γ RIIIa contains an ITAM motif and is therefore an activation receptor, whereas Fc γ RIIIb contains an ITIM motif and acts as an inhibitory

receptor [6]. In FcγRIIa, an important functional single nucleotide polymorphism (SNP) has been identified, resulting in an arginine (R) or a histidine (H) at amino acid position 131 [11]. These polymorphic variants have been shown to interact differently with IgG subclasses; FcγRIIa-H131 binds human IgG₂, whereas FcγRIIa-R131 does not [10,11]. A polymorphism in the transmembrane domain of FcγRIIb has been identified to affect the inhibitory activity of the ITIM motif [12]. Genetic polymorphisms affecting IgG-subclass binding also exist in *FCGR3A* [13] and *FCGR3B* genes [14]. As a result of such SNPs, the FcγRIIIa-V158 variant has a higher affinity for IgG₁ and IgG₃ than the FcγRIIIa-F158 variant [13]. Allelic variation in FcγRIIIb is comprised of differences in four amino acids, referred to as neutrophil antigen 1 (NA1) and NA2 [13]. FcγRIIIb-NA1 internalizes IgG₁- or IgG₃-opsonized particles more efficiently than FcγRIIIb-NA2 [14].

During recent years an increasing number of papers has been published on the linkage of various polymorphisms in the *FCGR* genes and a variety of diseases [15–25]. These diseases belong to the groups of infectious as well as autoimmune diseases. As it is generally accepted that Kawasaki disease is triggered by an infectious agent in genetically predisposed children, we investigated the possible association of functionally relevant SNPs in the *FCGR2* and *FCGR3* genes and Kawasaki disease. We hypothesized that this association could be true either for the entire cohort of Kawasaki disease patients (certain polymorphisms as a risk factor for increased susceptibility for Kawasaki disease) or for a subgroup of patients (as a risk factor for coronary artery damage and therapy failure).

Subjects and methods

Study population

In this study we included 167 patients with Kawasaki disease from our region, using the criteria for diagnosis and echocardiographic scoring according to the guidelines of the Kawasaki Disease Research Committee in Japan from 1984. Coronary artery dilatations with a diameter > 3 mm or > 1.5 times the adjacent vessel diameter were scored as aneurysmatic lesions. Standard therapy consisted of a single IVIG infusion (2 g/kg in 8–12 h) in combination with oral aspirin (80–100 mg/kg divided in four equal doses). A second IVIG infusion was administered when clinical symptoms (i.e. fever) persisted for 72 h or when echocardiographic findings were compatible with a progression or recurrence of CAL or the presence of a giant aneurysm (diameter > 8 mm). Echocardiographic findings were recorded before or shortly after the first IVIG infusion; follow-up was at 1–3-week intervals during the first 2–3 months, with an increase in frequency of visits depending on the presence, development or persistence of CAL. Healthy Caucasian blood donors (*n* = 239) were recruited

as controls. The controls were unrelated and lived in the same region as the Caucasian Kawasaki disease patients included in this study. Informed consent was obtained from all participants or parents/caregivers of the participants in the study. The study was approved by the medical ethical committee of the Academic Medical Centre (AMC), Amsterdam, the Netherlands.

Genotyping

We isolated the DNA from blood using the Puregene DNA isolation kit (Biozym; Hess, Oldendorf, Germany).

Primers and conditions for the polymerase chain reactions (PCR) for assessing the *FCGR* polymorphisms

The primers for the PCR reactions used to assess the *FCGR* polymorphisms are listed in Table 1. *FCGR2B* genotyping for the FcγRIIb-232I/T isoforms was performed by direct sequencing, using the same 2B/2C intron 4 primer used in PCR and an internal 2B/2C intron 5 reverse primer. The PCR condition for *FCGR2B* is as follows: the target DNA was amplified by 40 cycles of 94°C for 30 s, 60°C for 45 s and 72°C for 45 s, stimulated and heated lid (standardized condition in the DNA amplification of the PCR-product) at 105°C; for *FCGR2A* we used an allele-specific PCR with two internal control primers (Table 2), with conditions as follows: one cycle of 5 min 95°C, 2 min 58°C and 1 min 72°C, thereafter 10 cycles of 1 min 95°C, 2 min 58°C, 1 min 72°C and a final extension step for 5 min at 72°C. For *FCGR3A* the target DNA was amplified by one cycle at 95°C for 5 min, at 56°C for 1 min and at 72°C for 1 min, 30 cycles at 95°C for 1 min, at 56°C for 1 min and at 72°C for 1 min, and an elongation step at 72°C for 5 min. Direct sequencing was performed on the PCR product, using the same primers. For *FCGR3B* we used an allele-specific PCR with two internal control primers (Table 2). The amplification started with one cycle of 5 min at 95°C (denaturation), 1.5 min at 60°C (annealing), 2.5 min at 72°C (extension) and was followed by 10 cycles in which the denaturation time was reduced to 1 min. Subsequently, 24 cycles were performed consisting of 1 min at 95°C, 1 min at 57°C and 1 min at 72°C and finally one cycle of 1 min at 95°C, 1 min at 57°C and 5 min at 72°C.

Statistical analysis

Differences between Kawasaki patients and healthy controls in the frequencies of the polymorphisms for the various *FCGR* genotypes as well as of all possible combinations of the subtypes, were tested with χ^2 tests. The relationship between the *FCGR* subtypes as well as several known risk factors and development of CAL (yes/no) was studied with logistic regression analysis. All variables associated

Table 1. Primers for FCGR genotyping.

Primer name	Primer sequence
2B/2C intron 4 forward primer	5'-TGG GAC AAG GAG AGT ACT GCC TGT C-3'
2B intron 6 reverse primer	5'-TTT GAG CCC CAG CCA TCC TCC CAC T-3'
2B/2C intron 5 reverse primer	5'-GAA TGT GTA TCT AGC CCA AAG AGA G-3'
FCGR2A H131 forward primer	5'-ATC CCA GAA ATT CTC CCA-3'
FCGR2A R131 forward primer	5'-ATC CCA GAA ATT CTC CCG-3'
FCGR2A common reverse primer	5'-CAA TTT TGC TGC TAT GGG C-3'
HGH-I internal control primer	5'-CAG TGC CTT CCC AAC CAT TCC CTT A-3'
HGH-II internal control primer	5'-ATC CAC TCA CGG ATT TCT GTT GTG TTT C-3'
FCGR3A forward primer	5'-ATA TTT ACA GAA TGG CAC AGG-3'
FCGR3A reverse primer	5'-ACG TGC TGA GCT TGA GTG ATG ATG TTC AC-3'
FCGR3B NA1 forward primer	5'-CAG TGG TTT CAC GTG AA-3'
FCGR3B NA1 reverse primer	5'-ATG GAC TTC TAG CTG CAC CG-3'
FCGR3B NA2 forward primer	5'-CTC AAT GGT ACA GCG TGC TT-3'
FCGR3B NA2 reverse primer	5'-CTG TAC TCT CCA CTG TCG TT-3'
2221 internal control primer	5'-CTT GTG GGT AAA CCA AGG C-3'
2222 internal control primer	5'-TTT GGA AAA ACA CTG AGG TAA GTG GGG GT-3'

univariately with CAL with a P -value < 0.15 were entered in a multivariate logistic regression model to assess their independent prognostic value for CAL. The prognostic values of the variables were expressed as odds ratios (OR) with their 95% confidence intervals (95% CI). The OR can be interpreted as an estimation of the relative risk of CAL. A two-sided P -value < 0.05 was considered statistically significant. All analyses were performed with *SPSS* for Windows version 11.0 (SPSS Inc., Chicago, IL, USA).

Results

Baseline characteristics

Baseline characteristics are shown in Table 2. In this study 167 Caucasian Dutch children with Kawasaki disease were enrolled. Sufficient clinical and treatment data were available in 159 patients. The boy/girl ratio in this study was 1.8 : 1, which is comparable with the distribution in literature. The median age was 1 year and 10 months, with a range of 32 days–13.5 years. The standard treatment of 2 g/kg IVIG was administered in 149 of the patients (89%). Because of insufficient responsiveness to IVIG, 12 patients (7%) received corticosteroids as additional treatment. In 46 patients (28%) CAL developed during the course of the

disease; in eight patients these progressed to giant CAL (diameter > 8 mm). Two patients in our cohort died of the complications of Kawasaki disease.

Genotype and allelic frequencies

The frequency of the polymorphisms for the various *FCGR* subtypes were compared between children with Kawasaki disease and a healthy control population (Table 3). No difference was found in the distribution of any of the subtypes, nor in the allelic frequencies, when compared to the healthy controls. In addition, all possible combinations of the subtypes were analysed. Again, no difference was found in the distribution of any of the subtypes.

Table 2. Baseline characteristics.

	No. of patients	
	No.	(%)
Age, median (range)	1 year 10 months (32 days–13.5 years)	
Boys	107	(64%)
CAL	46	(28%)
Giant CAL	8	(5%)

CAL: coronary artery lesions.

Table 3. *FCGR* genotype frequencies in Kawasaki disease (KD) patients and healthy controls.

Genotype frequency	KD patients	Controls	χ^2	P -value
FcγRIIa	$n = 176$	$n = 239$		
131H/H	55 31 (%)	58 24 (%)	2.65	0.27
131H/R	92 52 (%)	134 56 (%)		
131R/R	29 17 (%)	47 20 (%)		
FcγRIIb	$n = 150$	$n = 239$		
232I/I	122 81 (%)	201 84 (%)	1.05	0.60
232I/T	27 18 (%)	35 15 (%)		
232T/T	1 1 (%)	3 1 (%)		
FcγRIIIa	$n = 172$	$n = 239$		
158F/F	68 40 (%)	99 41 (%)	0.36	0.84
158 V/F	83 48 (%)	115 48 (%)		
158 V/V	21 12 (%)	25 11 (%)		
FcγRIIIb	$n = 176$	$n = 239$		
NA1/NA1	34 19 (%)	34 14 (%)	1.92	0.38
NA1/NA2	80 46 (%)	115 48 (%)		
NA2/NA2	62 35 (%)	90 38 (%)		

Table 4. Univariate analysis for potential risk factors for developing coronary artery lesions in Kawasaki disease.

Factor	n	CAL	OR (95% CI)	P-value
FcγRIIa				
131H/H	43	12	Ref	
131H/R	75	21	1.01 (0.44–2.32)	0.990
131R/R	24	8	1.29 (0.44–3.80)	0.640
FcγRIIb				
232I/I	100	27	Ref	
232I/T	21	8	1.66 (0.62–4.46)	0.310
232T/T	1	0		
FcγRIIIa				
158F/F	55	17	Ref	
158 V/F	72	19	0.80 (0.37–1.74)	0.580
158 V/V	16	4	0.75 (0.21–2.65)	0.650
FcγRIIIb				
NA1/NA1	29	12	Ref	
NA1/NA2	66	15	0.42 (0.16–1.06)	0.067
NA2/NA2	49	13	0.51 (0.19–1.36)	0.180
Age				
< 1 year	51	23	Ref	
≥ 1 year	104	23	0.35 (0.17–0.71)	0.004
Treatment				
< 10 days	92	21	Ref	
≥ 10 days	44	20	2.82 (1.31–6.07)	0.008

Polymorphisms and other risk factors in relation to CAL

In Table 4 the univariate ORs of CAL associated with the polymorphisms for the *FCGR* genes and two known risk factors are presented. It was shown that age < 1 year and late onset of treatment were statistically significantly associated with an increased risk of CAL. From the *FCGR* genes tested, the *FCGR3B* genotype had a borderline statistically significant influence on the development of CAL. Here the *FCGR3B* genotypes for FcγRIIIb-NA1/NA2 had a slightly decreased risk of CAL compared to those for FcγRIIIb-NA1/NA1. After introduction of the most important univariate associations ($P < 0.15$) in a multivariate logistic regression analysis, only the variables age and onset of treatment remained statistically significantly independent risk factors for the development of CAL (age < 1 year: OR 0.36, 95% CI 0.14–0.88; treatment at > 10 days after onset of disease: OR 3.32, 95% CI 1.34–8.19). Although the *FCGR3b* genotype did not show a significant OR, a trend towards a protective effect was observed (Table 5). No interaction was observed between *FCGR3b* and onset of treatment on CAL.

Discussion

In this study we evaluated the hypothesis that SNPs for the genes coding for FcγRIIa, IIB, IIIa or IIIb are factors in the susceptibility or pathogenesis in Kawasaki disease. We com-

pared their frequencies because they all have influence either on the binding affinity for the IgG-isotypes of the various activating FcγRs (*FCGR2A*, *FCGR3A* and *FCGR3B*), or on the function of the inhibitory FcγR receptor (*FCGR2B*). We compared the distribution of these SNPs in a cohort of patients with Kawasaki disease with the distribution in a group of healthy Dutch Caucasian controls. It appeared that the distribution of the several allotypes was similar in the patient group when compared to a control population. Besides a borderline significant association for *FCGR3B*, no correlation between SNPs in FcγR genes and the development of CAL was found. The effect of *FCGR3B* on the risk of CAL appeared not to be confounded by the effect of age and late onset of treatment, as the point estimate of its OR and its 95% CI hardly changed after adjustment for both these variables in a multivariate regression model. Insufficient statistical power may be an explanation for lack of statistical significance. Therefore, an association between *FCGR3B* and CAL is not necessarily excluded by these results and needs to be explored further in larger numbers.

A recent study described a relationship between the FcγRIIa-131H/R genotype and the coronary outcome in 54 Japanese patients with Kawasaki disease [26]. Racial differences among Japanese and Dutch individuals are known to impact the allele frequency of some of the SNPs in the FcγRs [27]. This could explain the differences between Taniuchi and colleagues' study and ours, although we have tested 59 Japanese patients and 100 controls without finding any change in allele frequency between these two groups (data not shown).

Another study on FcγRs in Kawasaki disease described the regulation of receptor expression on circulating phagocytes in Kawasaki disease [28]. In this study the authors demonstrated an up-regulation of FcγRI and down-regulation of FcγRIII on neutrophils, and concomitant up-regulation of FcγRIII on monocytes. The authors concluded that the expression of FcγRs on circulating phagocytes could be an inflammatory marker of Kawasaki disease. In this study we analysed the association of single SNPs of various well-characterized and functionally relevant SNPs in the genes for

Table 5. Multivariate analysis for risk factors for developing coronary artery lesions in Kawasaki disease.

Factor	n	OR (95% CI)	P-value
FcγRIIIb			
NA1/NA1	20	Ref	
NA1/NA2	48	0.39 (0.12–1.27)	0.120
NA2/NA2	42	0.57 (0.18–1.82)	0.340
Age			
< 1 year	39	Ref	
≥ 1 year	71	0.36 (0.14–0.88)	0.025
Treatment			
< 10 days	73	Ref	
≥ 10 days	37	3.32 (1.34–8.19)	0.009

FcγRs or possible combinations thereof, with Kawasaki disease or disease severity as defined by the presence or absence of CAL by echocardiography. These SNPs have been associated with disease susceptibility or severity in various infectious diseases caused by, for example, *Neisseria meningitidis* [14,29–31] or *Streptococcus pneumoniae* [15,32,33], as well as in autoimmune disease [17–22,24,25]. As the single inhibitory IgG receptor, the FcγRIIb could seriously impact immunity in this respect. The genotype for the FcγRIIb-232 I/T isoforms represents a SNP in the transmembrane region of the receptor which was found to be important for B cell signalling [12]. Some reports link this receptor to susceptibility to SLE [27], although others have not been able to confirm this in a different racial background [34]. In a recent study, the same *FCGR2B* genotype for FcγRIIb-232 I/T appeared to be a predicting variable for chronic disease in newly diagnosed idiopathic thrombocytopenic purpura in childhood [35]. In Wegener's granulomatosis and in anti-neutrophil cytoplasmic antibody (ANCA)-associated vasculitis, associations were also found between polymorphisms in the genes for FcγRs by some groups [22,36] but not by others [23,37].

Although we did not find sufficient evidence for a correlation between the polymorphisms of the genes coding for the several FcγRs and Kawasaki disease, it does not rule out that (one of) these receptors are involved at some point during the disease process or treatment of the disease. The ratio of activating and inhibitory FcγRs on a specific immune cell may be an important factor in the resulting outcome of an immune response, as reported in various experimental animal models in FcγRIIb^{-/-} knock-out mice on inflammation, autoimmunity and an infectious model of *S. pneumoniae* septicaemia [38–40]. The identification of SNPs was performed in genomic DNA. It is hence impossible to investigate the role of transcription or expression factors. Moreover, these factors may differ among the several immune cells involved. Assessing, for example, the expression of the activating and inhibitory FcγRs on the inflammatory cells during several stages of Kawasaki disease might lead to new insights in the role of those receptors in this disease. However, specific antibodies to investigate this issue unequivocally are missing to date and such studies will have to wait for their availability.

Acknowledgements

We thank Dr Ludo van der Pol from the Department of Neurology, University Medical Center Utrecht, the Netherlands, for his help in composing the control group, and Dr M. Masuda for her kind cooperation in the collection of DNA from Japanese patients and controls. Maarten Biezeveld is funded by the Netherlands Heart Foundation (NHS 99-189). The funding sources had no role in the study design, data collection, data analysis, data interpretation or writing of the report.

References

- 1 Kawasaki T, Kosaki F, Okawa S *et al.* A new infantile acute febrile mucocutaneous lymph node syndrome (MLNS) prevailing in Japan. *Pediatrics* 1974; **54**:271–6.
- 2 Kawasaki T. [Acute febrile mucocutaneous syndrome with lymphoid involvement with specific desquamation of the fingers and toes in children]. *Arerugi* 1967; **16**:178–222.
- 3 Burns JC, Glode MP. Kawasaki syndrome. *Lancet* 2004; **364**:533–44.
- 4 Newburger JW, Takahashi M, Beiser AS *et al.* A single intravenous infusion of gamma globulin as compared with four infusions in the treatment of acute Kawasaki syndrome. *N Engl J Med* 1991; **324**:1633–9.
- 5 Tse SM, Silverman ED, McCrindle BW *et al.* Early treatment with intravenous immunoglobulin in patients with Kawasaki disease. *J Pediatr* 2002; **140**:450–5.
- 6 Ravetch JV, Bolland S. IgG Fc receptors. *Annu Rev Immunol* 2001; **19**:275–90.
- 7 Deo YM, Graziano RF, Repp R, van de Winkel JG. Clinical significance of IgG Fc receptors and Fc gamma R-directed immunotherapies. *Immunol Today* 1997; **18**:127–35.
- 8 Galon J, Robertson MW, Galinha A *et al.* Affinity of the interaction between Fc gamma receptor type III (Fc gammaRIII) and monomeric human IgG subclasses. Role of Fc gammaRIII glycosylation. *Eur J Immunol* 1997; **27**:1928–32.
- 9 Maenaka K, van der Merwe PA, Stuart DI, Jones EY, Sondermann P. The human low affinity Fc gamma receptors IIa, IIb, and III bind IgG with fast kinetics and distinct thermodynamic properties. *J Biol Chem* 2001; **276**:44898–904.
- 10 Maxwell KF, Powell MS, Hulett MD *et al.* Crystal structure of the human leukocyte Fc receptor, Fc gammaRIIa. *Nat Struct Biol* 1999; **6**:437–42.
- 11 Warmerdam PA, van de Winkel JG, Vlug A, Westerdaal NA, Capel PJ. A single amino acid in the second Ig-like domain of the human Fc gamma receptor II is critical for human IgG2 binding. *J Immunol* 1991; **147**:1338–43.
- 12 Li X, Wu J, Carter RH, Edberg JC *et al.* A novel polymorphism in the Fc gamma receptor IIB (CD32B) transmembrane region alters receptor signaling. *Arthritis Rheum* 2003; **48**:3242–52.
- 13 Koene HR, Kleijer M, Algra J, von Roos D, dem Borne AE, de Haas M. Fc gammaRIIIa-158V/F polymorphism influences the binding of IgG by natural killer cell Fc gammaRIIIa, independently of the Fc gammaRIIIa-48L/R/H phenotype. *Blood* 1997; **90**:1109–14.
- 14 de Haas M, Kleijer M, van Zwieten R, von Roos D, dem Borne AE. Neutrophil Fc gamma RIIB deficiency, nature, and clinical consequences: a study of 21 individuals from 14 families. *Blood* 1995; **86**:2403–13.
- 15 Bredius RG, Derkx BH, Fijen CA *et al.* Fc gamma receptor IIa (CD32) polymorphism in fulminant meningococcal septic shock in children. *J Infect Dis* 1994; **170**:848–53.
- 16 Norris CF, Surrey S, Bunin GR, Schwartz E, Buchanan GR, McKenzie SE. Relationship between Fc receptor IIA polymorphism and infection in children with sickle cell disease. *J Pediatr* 1996; **128**:813–9.
- 17 Manger K, Repp R, Spriewald BM *et al.* Fc gamma receptor IIa polymorphism in Caucasian patients with systemic lupus erythematosus: association with clinical symptoms. *Arthritis Rheum* 1998; **41**:1181–9.

- 18 Karassa FB, Trikalinos TA, Ioannidis JP. Role of the Fc gamma receptor IIa polymorphism in susceptibility to systemic lupus erythematosus and lupus nephritis: a meta-analysis. *Fc gamma RIIa, SLE Meta-Analysis Investigators. Arthritis Rheum* 2002; **46**:1563–71.
- 19 Kyogoku C, Tsuchiya N, Wu H, Tsao BP, Tokunaga K. Association of Fc gamma receptor IIA, but not IIB and IIIA, polymorphisms with systemic lupus erythematosus: a family-based association study in Caucasians. *Arthritis Rheum* 2004; **50**:671–3.
- 20 Atsumi T, Caliz R, Amengual O, Khamashta MA, Hughes GR. Fc gamma receptor IIA H/R131 polymorphism in patients with antiphospholipid antibodies. *Thromb Haemost* 1998; **79**:924–7.
- 21 Carlsson LE, Santoso S, Baurichter G *et al.* Heparin-induced thrombocytopenia: new insights into the impact of the Fc gamma RIIa-R-H131 polymorphism. *Blood* 1998; **92**:1526–31.
- 22 Dijkstra HM, Scheepers RH, Oost WW *et al.* Fc gamma receptor polymorphisms in Wegener's granulomatosis: risk factors for disease relapse. *Arthritis Rheum* 1999; **42**:1823–7.
- 23 Tse WY, Abadeh S, McTiernan A, Jefferis R, Savage CO, Adu D. No association between neutrophil Fc gamma RIIa allelic polymorphism and anti-neutrophil cytoplasmic antibody (ANCA)-positive systemic vasculitis. *Clin Exp Immunol* 1999; **117**:198–205.
- 24 van der Pol WL, van den Berg LH, Scheepers RH *et al.* IgG receptor IIA alleles determine susceptibility and severity of Guillain-Barre syndrome. *Neurology* 2000; **54**:1661–5.
- 25 Brun JG, Madland TM, Vedeler CA. Immunoglobulin G fc-receptor (Fc gamma R) IIA IIIA, and IIIB polymorphisms related to disease severity in rheumatoid arthritis. *J Rheumatol* 2002; **29**:1135–40.
- 26 Taniuchi S, Masuda M, Teraguchi M *et al.* Polymorphisms of Fc gamma RIIa may affect the efficacy of gamma-globulin therapy in Kawasaki disease. *J Clin Immunol* 2005 July; **25**:309–13.
- 27 Kyogoku C, Dijkstra HM, Tsuchiya N *et al.* Fc gamma receptor gene polymorphisms in Japanese patients with systemic lupus erythematosus: contribution of FCGR2B to genetic susceptibility. *Arthritis Rheum* 2002; **46**:1242–54.
- 28 Nakatani K, Takeshita S, Tsujimoto H, Kawamura Y, Kawase H, Sekine I. Regulation of the expression of Fc gamma receptor on circulating neutrophils and monocytes in Kawasaki disease. *Clin Exp Immunol* 1999; **117**:418–22.
- 29 Platonov AE, Shipulin GA, Vershinina IV, Dankert J, van de Winkel JG, Kuijper EJ. Association of human Fc gamma RIIa (CD32) polymorphism with susceptibility to and severity of meningococcal disease. *Clin Infect Dis* 1998; **27**:746–50.
- 30 Fijen CA, Bredius RG, Kuijper EJ *et al.* The role of Fc gamma receptor polymorphisms and C3 in the immune defence against *Neisseria meningitidis* in complement-deficient individuals. *Clin Exp Immunol* 2000; **120**:338–45.
- 31 Domingo P, Muniz-Diaz E, Baraldes MA *et al.* Associations between Fc gamma receptor IIA polymorphisms and the risk and prognosis of meningococcal disease. *Am J Med* 2002; **112**:19–25.
- 32 Abadi J, Zhong Z, Dobroszycki J, Pirofski LA. Fc gamma RIIa polymorphism in human immunodeficiency virus-infected children with invasive pneumococcal disease. *Pediatr Res* 1997; **42**:259–62.
- 33 Yee AM, Phan HM, Zuniga R, Salmon JE, Musher DM. Association between Fc gamma RIIa-R131 allotype and bacteremic pneumococcal pneumonia. *Clin Infect Dis* 2000; **30**:25–8.
- 34 Magnusson V, Zuncic R, Odeberg J *et al.* Polymorphisms of the Fc gamma receptor type IIB gene are not associated with systemic lupus erythematosus in the Swedish population. *Arthritis Rheum* 2004; **50**:1348–50.
- 35 Bruin M, Bierings M, Uiterwaal C *et al.* Platelet count, previous infection and FCGR2B genotype predict development of chronic disease in newly diagnosed idiopathic thrombocytopenia in childhood: results of a prospective study. *Br J Haematol* 2004; **127**:561–7.
- 36 Wainstein E, Edberg J, Csernok E *et al.* Fc gamma RIIIB alleles predict renal dysfunction in Wegener's granulomatosis (WG). *Arthritis Rheum* 1996; **39**:S210.
- 37 Tse WY, Abadeh S, Jefferis R, Savage COS, Adu D. Neutrophil Fc gamma RIIIB allelic polymorphism in anti-neutrophil cytoplasmic antibody (ANCA)-positive systemic vasculitis. *Clin Exp Immunol* 2000; **119**:574–7.
- 38 Samuelsson A, Towers TL, Ravetch JV. Anti-inflammatory activity of IVIG mediated through the inhibitory Fc receptor. *Science* 2001; **291**:484–6.
- 39 Ravetch JV, Lanier LL. Immune inhibitory receptors. *Science* 2000; **290**:84–9.
- 40 Clatworthy MR, Smith KGC. Fc gamma RIIb balances efficient pathogen clearance and the cytokine-mediated consequences of sepsis. *J Exp Med* 2004; **199**:717–23.